


I. Defining the fundamental group

Recall (X, p) based space


$f: [0,1] \rightarrow X$ loop $f(0) = f(1) = p$

$F: [0,1]^2 \rightarrow X$ homotopy between f_0, f_1 loops if

$F(s, i) = f_i$ for $i = 0, 1$, $F(0, t) = p = F(1, t)$


 $f_t := f(-, t): [0,1] \rightarrow X$ loop
vary t gives "path in the space of loops" from f_0 to f_1 .

write $f_0 \sim f_1$ homotopic.



Examples $X = \mathbb{R}^2$, $Y = \mathbb{R}^2 \setminus \{(0,0)\}$, $Z = \mathbb{R}^2 \setminus \{(1,0), (0,1)\}$

$$p = (1,0)$$


(a) Any loop $f: [0,1] \rightarrow X$ homotopic to a constant

("is nullhomotopic") via "straight-line homotopy"


$$F(s,t) = (1-t)f(s) + tp$$

(b) $g: [0,1] \rightarrow Y$ $g(s) = (\cos 2\pi s, \sin 2\pi s)$

g not homotopic to constant in Y .



(c) Z has many loops, distinct up to homotopy



Defined $\pi_1(X, p) = \{ \text{loops } f: [0, 1] \rightarrow X \text{ based at } p \} / \sim$

Denote $[f]$ equivalence class of f

Concatenation:



$$\pi_1(X, P) \times \pi_1(X, P) \longrightarrow \pi_1(X, P)$$

$$[f] \cdot [g] := [f * g]$$

$$(f * g)(s) = \begin{cases} f(2s) & 0 \leq s \leq \frac{1}{2} \\ g(2s - 1) & \frac{1}{2} \leq s \leq 1. \end{cases}$$

Well-defined

: if $f_i \sim f$ & $g_i \sim g$ then $f_i * g_i \sim f * g$.



$$H(s, t) = \begin{cases} F(2s, t) & s \in [0, \frac{1}{2}] \\ G(2s - 1, t) & s \in [\frac{1}{2}, 1] \end{cases}$$

$F * G$ ✓

Thm Concatenation of homotopy classes

Satisfies : for f, g, h any loops, $c \equiv p$ constant

(i) associativity

$$([f] \cdot [g]) \cdot [h] = [f] \cdot ([g] \cdot [h])$$


(ii) identity

$$[f] \cdot [c] = [f] = [c] \cdot [f]$$

(iii) inverses

$$[f] [\bar{f}] = [c] = [\bar{f}] [f]$$

where $\bar{f}(s) = f(1-s)$ reverse of f .



Cor $\pi_1(X, p)$ is a group under concatenation

Rmk This isn't true on level of maps


- need to consider homotopy classes.

$$f * c \neq f \quad (f * c)(s) = \begin{cases} f(2s) & 0 \leq s \leq \frac{1}{2} \\ p & \frac{1}{2} \leq s \leq 1. \end{cases} \neq f(s)$$

Proof of Thm

- identity

$$[f] * [c] = [f]$$




$$(f * c)(s) = \begin{cases} f(2s) & 0 \leq s \leq \frac{1}{2} \\ P & \frac{1}{2} \leq s \leq 1 \end{cases}$$

$$[2(1-t) + t \cdot 1] = 2 - 2t + t = 2 - t$$

$$F(s, t) = \begin{cases} f((2-t) \cdot s) & 0 \leq s \leq \frac{1}{2-t} \\ P & \frac{1}{2-t} \leq s \leq 1 \end{cases}$$

✓


- Inverses



$$(f * \bar{f})(s) = \begin{cases} f(2s) & 0 \leq s \leq \frac{1}{2}, \\ \bar{f}(2s-1) & \frac{1}{2} \leq s \leq 1 \end{cases}$$

$$= \begin{cases} f(\underline{2s}) & 0 \leq s \leq \frac{1}{2}, \\ f(\underline{2-2s}) & \frac{1}{2} \leq s \leq 1 \end{cases}$$

$$F(s,t) = \begin{cases} f(2ts) & 0 \leq s \leq \frac{1}{2}, \\ f(2t(1-s)) & \frac{1}{2} \leq s \leq 1. \end{cases}$$




• associativity : you

$$(f * g) * h$$

$$f * (g * h)$$

$$\begin{array}{c} \hline | & | & | \\ f & g & h \end{array}$$

$$\begin{array}{c} \hline | & | & | \\ f & g & h \end{array}$$



□