


I. Computing edge groups

Recap K simplicial complex $K \subset P(V)$


eg graphs: V vertex set $G \subset P(V)$ subsets of size $e \leq 2$.



given K , p vertex \rightsquigarrow edge group

$$E(K, p) = \{ \text{edge loops at } p \} / \sim$$

Eg $V = \{0, 1, 2\}$ $K = P(V) \setminus \emptyset$



$$E(K, p) \supseteq [0120], [0210], [01010].$$

$$0120 \sim 020 \sim 0.$$

these are all equivalent to $[0]$ eg

Give a presentation for $E(K, \rho)$.

Assume K connected (any $u, v \in V$ are connected
by an edge path)

Write $V = \{v_i\}$


Choose max tree $L \subset K$ (cf. HW 1 # 1)

Define $G(K, L)$ group with

- generators g_{ij} for each edge $\{v_i, v_j\}$ $i < j$
- relations
 - $g_{ij} g_{ik} = g_{ik}$ if $\{v_i, v_j, v_k\} \in K$ $i < j < k$
 - $g_{ij} = 1$ if $\{v_i, v_j\} \in L$.

Thm $G(K, L) \cong E(K, p)$

Ex.



generators: $g_{01}, g_{03}, g_{12}, g_{13}, g_{23}$

relations: $g_{12}g_{23} = g_{13}$

$$G(K, L) = \langle g_{01} = 1, g_{03} = 1, g_{23} = 1 \rangle$$

$$\langle \cancel{g_{01}}, \cancel{g_{03}}, g_{12}, g_{13}, \cancel{g_{23}} \mid g_{01} = g_{03} = g_{23} = 1, g_{12}\cancel{g_{23}} = g_{13} \rangle$$

$$\cong \langle g_{12} \mid \rangle \cong \mathbb{Z}$$


Remark This gives reasonable way to compute

$\pi_1(X)$ if X has a triangulation $X \cong |K|$

Since $\pi_1(X) \cong \pi_1(|K|) \cong E(K, \rho) \cong G(K, L)$


need proof

Ex $|K| \cong S^1$



$$G(K, L) = \langle g_{12}, g_{23}, g_{31} \mid g_{13} = 1 = g_{23} \rangle$$
$$= \langle g_{12} \rangle \cong \mathbb{Z}$$

Ex $|K| \cong S^2$



$$G(K, L) = \langle 1 \rangle = \{1\}.$$


generators g_{12} g_{13} g_{14} g_{23} g_{24} g_{34}

relations $g_{12}g_{23} = g_{13}$ $g_{12}g_{24} = g_{14}$

$$g_{13}g_{34} = g_{14} \quad g_{23}g_{34} = g_{24}$$

$$g_{14} = 1 \quad g_{24} = 1 \quad g_{34} = 1$$

Ex $|K| \cong S' \vee S'$




$$G(K, L) = \langle g_{12}, g_{34} \rangle$$

generators $g_{12}, g_{\cancel{01}}, g_{\cancel{02}}, g_{\cancel{03}}, g_{\cancel{04}}, g_{34}$

relations $g_{01} = 1 = g_{02} = g_{03} = g_{04}$

a group with no relations is called a free group.

$$\langle a, b \rangle \ni a b a^{-1} b^2 a^{-7} b$$



Remarks

- $G(K, L)$ doesn't use any info about simplices of K of $\dim \geq 3$ (only use vertices, edges, faces)
- it can be tedious to write down $G(K, L)$ if K has many simplices. The presentation for $G(K, L)$ is often very redundant.
- This is problematic: \nexists algorithm to decide if two presentations $\langle S | R \rangle$ and $\langle S' | R' \rangle$ define isomorphic groups. eg can't decide in general if $\langle S | R \rangle = \{1\}$.

• For different choices of L, K get different

presentations: $G(K, L) \stackrel{?}{\cong} G(K, L')$

Thm $E(K, P) \cong G(K, L)$